

Vaccination Program

Foals (based off age)

3 months – Botulism (there is no maternal antibody interference with the Botulism vaccine)

4 months – EWT + West Nile, Botulism, Flu/Rhino

5 months – EWT + West Nile, Botulism, Flu/Rhino (PHF if desired)

6 months – EWT + West Nile, Rabies (PHF if desired)

7 months – Rabies, Flu/Rhino

- If dam was unvaccinated, Rabies is instead administered at 3 and 4 months. Also, EWT/FR + West Nile series may be given at an earlier age. *Please consult with your veterinarian*
- If vaccinating for Strangles (intranasal):
 - If foal is at risk – can begin series of 3 doses given 4 weeks apart starting as early as 6 weeks of age, although recommended to begin at 6-9 months of age.

If vaccinating with Flu (intranasal) – give first dose at 6 months of age and second dose at 11 months of age

Yearlings

Vaccinated patients:

***Vaccinate all yearlings in January regardless of age

January – Rabies, Flu/Rhino

April – Flu/Rhino booster, West Nile, EWT, Botulism

July – Flu/Rhino (PHF if desired)

October – Flu/Rhino

Strangles intranasal and Flu intranasal require annual boosters

Unvaccinated patients:

EWT + WNV: 2 dose series, given 4 weeks apart

Rabies: 1 dose series

Botulism: 3 dose series, given 4 weeks apart

Flu/Rhino: 3 dose series, given 4 weeks apart

PHF: 2 dose series, given 4 weeks apart

Strangles intranasal: 2 dose series, given 4 weeks apart

Flu intranasal: 1 dose series

Adults

Vaccinated patients:

All vaccines require an annual booster (*exception: Flu/Rhino should be given every 6 months*)

Unvaccinated patients:

EWT + WNV: 2 dose series, given 4 weeks apart

Rabies: 1 dose series

Botulism: 3 dose series, given 4 weeks apart

Flu/Rhino: 3 dose series, given 4 weeks apart

PHF: 2 dose series, given 4 weeks apart

Strangles intranasal: 2 dose series, given 4 weeks apart

Flu intranasal: 1 dose series

Pregnant Mares

Observe all vaccination recommendations for adult horses with the following modifications:

- 5 months – EHV
- 7 months – EHV
- 8 months – Rota
- 9 months – EHV, Rota
- 10 months – EWT, West Nile, Flu/Rhino (Killed only!), Rabies, Botulism, Rota

Note:

- ***Horses leaving the farm for training or racing, breeding or sales should have a Rhino vaccine every 3 months***
- Breeding stallions should not be vaccinated during breeding season. NO vaccines should be given 60 days before the start of the season
 - Stallions must be vaccinated for EVA annually. They are in quarantine for 28 days following vaccination
 - A negative EVA must be recorded before vaccinating for EVA